






How to give the right amount of IBUPROFEN (also known as Motrin, Advil) is different depending on which type of ibuprofen you plan to give.

Dose: Give every 6 hours if needed, for fever or pain. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.
 Do NOT use with any other medicine containing ibuprofen.

Weight	Age	Infant's Ibuprofen Drops (50 mg / 1.25 mL)	Children's Liquid Ibuprofen (100 mg / 5 mL)	Children's Ibuprofen Chewable Tablets (50 mg)	Junior Strength Ibuprofen Tablets (100 mg)	Adult's Ibuprofen Tablets (200 mg)
0 to 11 pounds (up to 5 kilograms)	0 to 5 months	_____	_____	_____	_____	_____
12 to 17 pounds (about 6 to 7 kilograms)	6 to 11 months	1.25 mL 	2.5 mL 	_____	_____	_____
18 to 23 pounds (about 8 to 10 kilograms)	12 to 23 months	1.875 mL 	4 mL 	_____	_____	_____
24 to 35 pounds (about 11 to 15 kilograms)	2 to 3 years	2.5 mL 	5 mL 	2 tablets 	_____	_____
36 to 47 pounds (about 16 to 21 kilograms)	4 to 5 years	3.75 mL 	7.5 mL 	3 tablets 	_____	_____
48 to 59 pounds (about 22 to 26 kilograms)	6 to 8 years	5 mL 	10 mL 	4 tablets 	2 tablets 	1 tablet 
60 to 71 pounds (about 27 to 32 kilograms)	9 to 10 years	_____	12.5 mL 	5 tablets 	2 1/2 tablets 	1 tablet 
72 to 95 pounds (about 33 to 43 kilograms)	11 years	_____	15 mL 	6 tablets 	3 tablets 	1 1/2 tablets 
96 pounds or more (44 kilograms or more)	12 years or older	_____	10 mL  and 10 mL  10 mL + 10 mL = 20 mL	8 tablets 	4 tablets 	2 tablets 